

Deficiency of LKB1 in skeletal muscle prevents AMPK activation and glucose uptake during contraction

Kei Sakamoto^{1,*}, Afshan McCarthy²,
Darrin Smith², Kevin A Green³,
D Grahame Hardie³, Alan Ashworth²
and Dario R Alessi¹

¹MRC Protein Phosphorylation Unit, University of Dundee, Dundee, UK,

²The Breakthrough Breast Cancer Research Centre, The Institute of Cancer Research, London, UK and ³Division of Molecular Physiology, School of Life Sciences, University of Dundee, Dundee, UK

Recent studies indicate that the LKB1 tumour suppressor protein kinase is the major 'upstream' activator of the energy sensor AMP-activated protein kinase (AMPK). We have used mice in which LKB1 is expressed at only ~10% of the normal levels in muscle and most other tissues, or that lack LKB1 entirely in skeletal muscle. Muscle expressing only 10% of the normal level of LKB1 had significantly reduced phosphorylation and activation of AMPK α 2. In LKB1-lacking muscle, the basal activity of the AMPK α 2 isoform was greatly reduced and was not increased by the AMP-mimetic agent, 5-aminoimidazole-4-carboxamide riboside (AICAR), by the antidiabetic drug phenformin, or by muscle contraction. Moreover, phosphorylation of acetyl CoA carboxylase-2, a downstream target of AMPK, was profoundly reduced. Glucose uptake stimulated by AICAR or muscle contraction, but not by insulin, was inhibited in the absence of LKB1. Contraction increased the AMP:ATP ratio to a greater extent in LKB1-deficient muscles than in LKB1-expressing muscles. These studies establish the importance of LKB1 in regulating AMPK activity and cellular energy levels in response to contraction and phenformin.

The EMBO Journal (2005) 24, 1810–1820. doi:10.1038/sj.emboj.7600667; Published online 5 May 2005

Subject Categories: signal transduction; cellular metabolism

Keywords: AMP-activated protein kinase; glucose transport; LKB1; phenformin; skeletal muscle

Introduction

The AMP-activated protein kinase (AMPK) is a major sensor of cellular energy levels and is activated by increased levels of 5'-AMP resulting from reduced energy availability (reviewed in Carling, 2004; Hardie, 2004). AMPK is a heterotrimeric complex formed from a catalytic α subunit and two regulatory subunits termed AMPK β and AMPK γ . AMP activates the

AMPK complex by binding to two sites formed by the CBS motifs located on the γ subunit (Scott *et al*, 2004). Once activated, AMPK phosphorylates substrates that inhibit anabolic processes and promote catabolic processes to restore cellular energy levels. One of the most studied physiological events that activates AMPK is exercise/contraction in skeletal muscle, where AMPK has been proposed to play a key role in stimulation of glucose uptake. Thus, treatment of muscle with an AMPK activator, 5-aminoimidazole-4-carboxamide riboside (AICAR), which is converted to an AMP mimetic within the cell, stimulated glucose uptake (Merrill *et al*, 1997; Hayashi *et al*, 1998). Moreover, overexpression of a constitutively active AMPK mutant in muscle cell lines increased glucose uptake (Fryer *et al*, 2002), while overexpression of a dominant-negative mutant of AMPK in mouse skeletal muscle blocked AICAR-induced glucose uptake (Mu *et al*, 2001). However, glucose uptake induced by contraction was only inhibited partially in mouse muscle expressing dominant-negative AMPK (Mu *et al*, 2001), suggesting that AMPK-independent pathway(s) may regulate glucose transport in contracting muscle. AMPK is also activated by metformin, the most widely utilised drug for reducing blood glucose levels in Type II diabetics (Zhou *et al*, 2001). The mechanism by which metformin, or its closely related analogue phenformin, activates AMPK has not yet been fully established, but may involve inhibition of ATP production via effects on complex I of the mitochondrial respiratory chain.

Activation of AMPK by AICAR, metformin and contraction requires phosphorylation of a threonine residue in the T-loop of the AMPK α subunit kinase domain (corresponding to Thr172 in both the AMPK α 1 and AMPK α 2 isoforms), by upstream kinase(s) (Carling, 2004; Hardie, 2004). Work performed initially in *Saccharomyces cerevisiae* (Hong *et al*, 2003; Nath *et al*, 2003; Sutherland *et al*, 2003), and subsequently in mammalian cells (Hawley *et al*, 2003; Woods *et al*, 2003a; Shaw *et al*, 2004), demonstrated that the LKB1 protein kinase can mediate Thr172 phosphorylation of AMPK both *in vitro* and in intact cells. LKB1 is a 50 kDa serine/threonine kinase that was originally identified as the product of the gene mutated in the autosomal dominantly-inherited Peutz-Jeghers cancer syndrome (Hemminki *et al*, 1998; Jenne *et al*, 1998). Like AMPK, LKB1 forms a heterotrimeric complex, in this case with regulatory proteins termed STRAD and MO25, which are required for its activation and cytosolic localisation (Baas *et al*, 2003; Boudeau *et al*, 2003, 2004). The LKB1 complex is not itself stimulated by AMP, and is constitutively active in cell lines (Woods *et al*, 2003a; Lizcano *et al*, 2004), as well as in skeletal muscle (Sakamoto *et al*, 2004). *In vitro* studies have suggested that binding of AMP to AMPK is likely to be the principle regulatory mechanism stimulating phosphorylation of AMPK by LKB1 (Hawley *et al*, 2003). The LKB1 complex also phosphorylates and activates a number of other protein kinases related to AMPK, whose functions are poorly understood (Lizcano *et al*, 2004; Jaleel *et al*, 2005).

*Corresponding author. MRC Protein Phosphorylation Unit, School of Life Sciences, University of Dundee, Dow Street, Dundee DD1 5EH, UK. Tel.: +44 1382 34 4241; Fax: +44 1382 223 778; E-mail: k.sakamoto@dundee.ac.uk

Received: 22 December 2004; accepted: 31 March 2005; published online: 5 May 2005

However, unlike AMPK, the other AMPK-related kinases phosphorylated by LKB1 are not activated by AICAR, phenformin or muscle contraction (Lizcano *et al*, 2004; Sakamoto *et al*, 2004).

Recent studies have suggested that activation of AMPK might be mediated independently of LKB1 in IGF1-stimulated fibroblasts and HeLa cells (Suzuki *et al*, 2004) or in cardiac muscle in response to ischaemia (Altarejos *et al*, 2005; Baron *et al*, 2005). Furthermore, in LKB1-deficient HeLa cells or mouse embryo fibroblasts, although AMPK is not activated by AICAR or other stimuli tested, it did possess significant basal activity and phosphorylation at Thr172 (Hawley *et al*, 2003; Shaw *et al*, 2004). This suggested that AMPK could be phosphorylated at Thr172 *in vivo* independently of LKB1. In this study, we wished to address the role that LKB1 plays in activating AMPK in a mammalian tissue *in vivo*, rather than in cultured cells *in vitro*. We have generated mice either lacking LKB1 in muscle, or in which the expression of LKB1 is reduced 10-fold, to define the role that LKB1 plays in regulating AMPK activity and glucose uptake in contracting skeletal muscle.

Results

Strategy employed to generate muscle-specific LKB1 knockout mice

We have generated mice conditional for expression of LKB1 and the structure of the floxed (fl) allele is shown in Figure 1A. Exons 5–7 of the *LKB1* gene were replaced by a cDNA cassette encoding the remainder of the LKB1 sequence, and exons 4 of the *LKB1* gene encoding residues 156–203 and the cDNA cassette are flanked by the *loxP* Cre excision sequence. This strategy was employed to ensure that in the *LKB1^{fl/fl}* mice, full-length wild-type LKB1 would be expressed

after exon 4 from the LKB1 cDNA cassette. The *LKB1^{fl/fl}* mice were crossed with transgenic mice expressing the Cre recombinase under the muscle creatine kinase promoter, which induces expression of the Cre recombinase specifically in skeletal muscle and heart, just prior to birth (Bruning *et al*, 1998). In the resulting *LKB1^{fl/fl}Cre^{+/-}* mice, exon 4 as well as the LKB1 cDNA cassette would be specifically excised in skeletal and cardiac muscle, ablating functional LKB1 expression, but potentially resulting in expression of an N-terminal fragment of LKB1 encompassing exons 1–3 (encoding residues 1–155).

Hypomorphic phenotype of *LKB1^{fl/fl}* mice

LKB1^{fl/fl} mice were bred as described in Cross 1 in Figure 1B and we observed that these were born at ~30% lower than expected Mendelian frequency. Female *LKB1^{fl/fl}* mice were fertile (Cross 2, Figure 1B) but, unexpectedly, male *LKB1^{fl/fl}* mice were infertile (Cross 3, Figure 1B). To investigate whether this might result from reduced expression of LKB1, we quantified the level of LKB1 protein and activity in various tissues including testis, from 7–10-week-old *LKB1^{fl/fl}* animals (Figure 2). This revealed that in testis, as well as skeletal muscle, heart, liver and kidney, LKB1 protein and activity was five- to 10-fold lower in *LKB1^{fl/fl}* mice compared to littermate *LKB1^{+/+}* mice. The heterozygous *LKB1^{+/-}* mice expressed two-fold lower levels of LKB1 compared to *LKB1^{+/+}* mice. In testis, in addition to the 50 kDa protein corresponding to full-length LKB1, a major faster migrating ~48 kDa species was also recognised by the LKB1 antibody in an immunoblot analysis (Figure 2F). Interestingly, the faster migrating species was not detected in the *LKB1^{fl/fl}* mice, while the 50 kDa species was reduced ~10-fold. In contrast to other tissues, the levels of LKB1 in the brain of *LKB1^{fl/fl}* mice were only reduced slightly (Figure 2G).




Figure 1 Generation of LKB1-deficient mice. (A) Diagram illustrating the positions of exons 3–8 (■) in the wild-type *LKB1⁺Cre⁻* allele. In the *LKB1^{fl}Cre⁻* allele, exon 4 of the *LKB1* gene is flanked with *loxP* Cre recombinase excision sites (▲), and exons 5–7 encoding the catalytic domain of LKB1 are replaced with a cDNA construct encoding the remainder of the LKB1 sequence, as well as a neomycin (Neo) selection gene. The expression of the neomycin gene is driven by the LKB1 promoter and it is made as fusion mRNA with LKB1. Its translation is directed by an internal ribosome entry site. In the *LKB1^{fl}Cre⁺* allele, exons 4–7 of the *LKB1* gene are deleted through action of the Cre recombinase, thereby ablating functional expression of LKB1. The positions of the PCR primers used to genotype the mice are indicated with arrows. (B) Breeding strategy employed to generate *LKB1^{fl/fl}* and LKB1-muscle-deficient (*LKB1^{fl/fl}Cre^{+/-}*) mice, where MCKCre denotes transgenic mice expressing the Cre recombinase under the muscle creatine kinase promoter. The number and percentage of each genotype obtained are indicated.




Figure 2 Expression and activity of LKB1 in mouse tissues. (A–G) Extracts of the indicated muscles and other tissues derived from wild-type and mutant mice (7–10 weeks of age) were generated. LKB1 activity was assessed following its immunoprecipitation and assay with the LKBtide peptide. Assays were performed in duplicate from tissues derived from three mice and results presented as the average \pm s.e.m. The level of LKB1 protein was assessed by immunoblot analysis of 30 μ g of protein from tissue extracts using either anti-LKB1 antibody raised against the whole mouse LKB1 protein (WP antibody) or an anti-LKB1 N-terminal antibody raised against the peptide encompassing residues 24–39 of human LKB1 (NT antibody). The ERK1 and ERK2 kinases were immunoblotted as a loading control. The immunoblotting results are representative of independent experiments performed with tissues from two mice. (a) $P < 0.05$ versus LKB1^{+/+}; (b) $P < 0.05$ versus LKB1^{+/ Δ} Cre^{-/-}; (c) $P < 0.05$ versus LKB1 ^{Δ / Δ} Cre^{-/-}. Statistical analysis performed using unpaired Student's *t* test.

LKB1 expression is abolished in the muscle of LKB1 ^{Δ / Δ} Cre^{+/-} mice

We next analysed LKB1 in tissues of the LKB1 ^{Δ / Δ} mice expressing the Cre recombinase, generated as described in Cross 2 in Figure 1B. We employed antibodies raised against

the full-length LKB1 protein as well as an antibody raised against an N-terminal peptide to enable us to detect any expression of the N-terminal exon 1–3 fragment of LKB1, which would be expected to have a molecular mass of \sim 17 kDa. We found that, in the LKB1 ^{Δ / Δ} Cre^{+/-} mice, no

LKB1 expression or activity was detected using either antibody in quadriceps muscle (Figure 2A), indicating that Cre ablated expression of full-length LKB1 and that the exon 1–3 fragment of LKB1 is unstable. In three other skeletal muscle types or in cardiac muscle, expression of LKB1 was abolished in LKB1^{fl/fl}Cre^{+/-} animals (Figure 2B and C). As expected, in nonmuscle tissues of LKB1^{fl/fl}Cre^{+/-} mice, LKB1 was expressed at levels similar to those found in LKB1^{fl/fl}Cre^{-/-} littermate mice (Figure 2D–G). The LKB1^{fl/fl}Cre^{+/-} mice, as well as their hypomorphic littermate LKB1^{fl/fl}Cre^{-/-} mice, displayed no overt phenotype up to the age of 10 weeks. They possessed normal fasted and fed blood glucose levels, and growth curves from 4 to 10 weeks of age indicated that these animals were of normal size and weight compared to control littermates (data not shown).

LKB1 is required for AICAR-induced AMPK α 2 activation

To determine whether lack of LKB1 in skeletal muscle affected AMPK activation, we treated isolated extensor digitorum longus (EDL) muscle isolated from 7–10-week-old mice, in the presence or absence of 2 mM AICAR, which is converted by adenosine kinase into AICAR monophosphate (ZMP), a cellular mimetic of AMP which activates AMPK (Corton *et al*, 1995). We first measured the activity of the AMPK α 2 isoform as well as its phosphorylation at Thr172, the site of LKB1 phosphorylation. In LKB1^{+/+} and LKB1^{+/fl} EDL muscles, AICAR stimulated AMPK α 2 activity ~2-fold (Figure 3A) and robustly increased phosphorylation at Thr172 (Figure 3B). In contrast, in the EDL muscle of LKB1^{fl/fl}Cre^{+/-} lacking LKB1 activity, the basal AMPK α 2 activity (1.8 mU/mg) was much lower than that observed (~40 mU/mg) in LKB1-expressing LKB1^{+/+} or LKB1^{+/fl} muscle, although AMPK α 2 was expressed normally. Moreover, in LKB1-deficient muscle, AICAR barely increased AMPK α 2 activity (to 2.3 mU/mg) compared to ~80 mU/mg in LKB1-expressing LKB1^{+/+} or LKB1^{+/fl} muscle. Consistent with the lack of AMPK α 2 activation in LKB1^{fl/fl}Cre^{+/-} muscle, AICAR also failed to induce detectable phosphorylation of Thr172 (Figure 3B). AMPK activity *in vivo* was also assessed by analysing the phosphorylation of a downstream target of AMPK, namely the muscle isoform of acetyl-CoA carboxylase-2/ β (ACC-2/ β) isoform, at the primary site phosphorylated by AMPK (Ser212, equivalent to Ser79 phosphorylated on ACC-1/ α ; Davies *et al*, 1990). In LKB1^{+/+} or LKB1^{+/fl} muscle, significant basal levels of ACC phosphorylation were observed, which were profoundly increased by AICAR (Figure 3B). In contrast, in both unstimulated and AICAR-treated LKB1^{fl/fl}Cre^{+/-} muscle, phosphorylation of ACC was undetectable.

Further evidence that LKB1 regulated AMPK α 2 activity came from the finding that, in the LKB1^{fl/fl} hypomorphic mice, which express 10-fold lower levels of LKB1 in muscle (Figure 2A), the basal and AICAR-stimulated levels of AMPK α 2 activity and Thr172 phosphorylation were reduced from four- to two-fold, compared to LKB1^{+/+} and LKB1^{+/fl} muscle (Figure 3A and B). The phosphorylation of ACC in response to AICAR was reduced by 60% in the hypomorphic mice, as judged by quantitative immunoblot analysis using the Li-Cor Odyssey system (Figure 3B).

We next examined AMPK α 1 activity in the EDL muscle of wild-type LKB1^{+/+} mice and found that its activity was markedly lower than that of AMPK α 2 (compare Figure 3A

and C), although AICAR still stimulated AMPK α 1 activity two-fold. In the hypomorphic LKB1^{fl/fl}Cre^{-/-} mice, AMPK α 1 was expressed normally, but its basal and AICAR-stimulated activities were reduced ~4-fold compared to LKB1^{+/+} and LKB1^{+/fl} muscle, suggesting that LKB1 does regulate AMPK α 1 activity. In LKB1-deficient LKB1^{fl/fl}Cre^{+/-} muscle, we noticed that AMPK α 1 protein levels were increased ~2-fold, as judged by quantitative immunoblot analysis using the Li-Cor Odyssey system (Figure 3B). Interestingly however, the basal and stimulated AMPK α 1 activities were not further reduced in LKB1-deficient LKB1^{fl/fl}Cre^{+/-} compared with LKB1^{fl/fl}Cre^{-/-} muscle, and AMPK α 1 was still modestly activated by AICAR. This result might be explained if the AMPK α 1 activity was derived from a minor amount of nonmuscle cells in the preparation, which would not express the Cre recombinase.

LKB1 is required for AICAR-induced glucose transport in skeletal muscle

A major physiological role of AMPK in muscle is to stimulate glucose transport (Winder and Hardie, 1999). We therefore measured 2-deoxyglucose uptake in isolated EDL muscle incubated for 60 min in the presence or absence of 2 mM AICAR, followed by 10 min in ³H-2-deoxyglucose. These studies revealed that the basal levels of glucose uptake were similar in LKB1^{+/+} or hypomorphic LKB1^{fl/fl}Cre^{-/-} muscle, and that glucose uptake was stimulated two- to three-fold by AICAR (Figure 3D). In contrast, in the LKB1-deficient LKB1^{fl/fl}Cre^{+/-} muscle, the basal uptake of ³H-2-deoxyglucose was similar to that observed in LKB1-expressing muscle, but AICAR-stimulated glucose uptake was blocked completely (Figure 3D).

LKB1 is required for contraction-induced AMPK α 2 activation

One of the major physiological activators of AMPK in skeletal muscle is contraction or exercise (Winder and Hardie, 1999). In order to investigate the role of LKB1 in enabling AMPK to be activated by muscle contraction, *in situ* contractions of hind limb muscle were induced in one leg via electrical stimulation of the sciatic nerve in anaesthetised mice, while the other leg served as the noncontracted control. Contraction was performed for 5 min, a time point that maximally activates AMPK α 2 using the protocol employed (Sakamoto *et al*, 2004). We measured AMPK α 2 activity (Figure 4A), and phosphorylation of AMPK at Thr172 and ACC at Ser212, in mixed tibialis anterior and EDL muscles (Figure 4B). In LKB1^{+/+} and LKB1^{+/fl} muscle, contraction increased AMPK α 2 activity six-fold and also induced marked phosphorylation of AMPK at Thr172 and ACC at Ser212. In contrast, in the LKB1-deficient LKB1^{fl/fl}Cre^{+/-} muscle, the basal and contraction-induced AMPK α 2 activities were reduced ~50-fold, compared to LKB1^{+/+} and LKB1^{+/fl} muscles (Figure 4A). Furthermore, in LKB1-deficient muscle, contraction failed to stimulate phosphorylation of AMPK at Thr172, while phosphorylation of ACC was reduced by nearly 90% (Figure 4B).

In the hypomorphic LKB1^{fl/fl}Cre^{-/-} muscle, the basal AMPK α 2 activity was 6 mU/mg (compared to 25 mU/mg in LKB1^{+/+} muscle) and was stimulated during contraction to 55 mU/mg (compared to 115 mU/mg in LKB1^{+/+} muscle). In the LKB1 hypomorphic mice, contraction-induced phosphor-




Figure 3 Role of LKB1 in regulating AICAR-induced AMPK activation and glucose transport. (A) Isolated mouse EDL muscles derived from wild-type and mutant mice (7–10 weeks of age) were incubated in the presence (+) or absence (–) of 2 mM AICAR for 60 min. AMPK α 2 was immunoprecipitated and assayed with the AMARA peptide. Assays were performed in duplicate from tissues derived from 4–5 mice and results are presented as the average \pm s.e.m. (B) Equal amounts of protein (20–30 μ g) from muscle extracts were immunoblotted with the indicated antibodies. The immunoblotting results are representative of independent experiments performed with tissues from at least three mice. Immunoblot analysis of AMPK α 1 and AMPK α 2 levels as well as ACC phosphorylation was also assessed by quantitative Li-Cor analysis. The data presented are the mean \pm s.e.m. Expression relative to expression in LKB1^{+/+} muscle derived from 3–4 mice. (C) As in (A), except that AMPK α 1 was immunoprecipitated and assayed. (D) Glucose transport in isolated EDL muscle derived from the indicated mice was determined without (–) or with (+) 2 mM AICAR, as described in Materials and methods. The data are presented as the mean \pm s.e.m. for muscle isolated from 4–5 mice for each genotype. * P < 0.05 basal versus AICAR within each genotype; (a), P < 0.05 versus LKB1^{+/+} (basal); (b) P < 0.05 versus LKB1^{fl/fl}Cre^{-/-} (basal); (c) P < 0.05 versus LKB1^{+/+} (AICAR); (d) P < 0.05 versus LKB1^{fl/fl}Cre^{-/-} (AICAR). Statistical analysis performed using one-way ANOVA and Tukey's *post hoc* test.

ylation of AMPK at Thr172 was reduced significantly, but, in contrast to AICAR treatment (Figure 3B), the phosphorylation of ACC at Ser212 was similar to that observed in LKB1^{+/+} muscle (Figure 4B).

In contrast to AMPK α 2, AMPK α 1 activity was not increased significantly by muscle contraction (Figure 4C), consistent with the notion outlined above that AMPK α 1 detected in mixed tibialis anterior and EDL muscle may largely be derived from nonmuscle cells. Previous studies have also found that the AMPK α 1 is not activated in contracting rat or human muscle under conditions where AMPK α 2 is stimulated (Fujii *et al*, 2000). AMPK α 1 activity was \sim 3-fold lower in hypomorphic LKB1^{fl/fl}Cre^{-/-} muscle compared to LKB1^{+/+} muscle, and its activity was not reduced further in the

Cre-expressing muscles (Figure 4C; see also Figure 3C). In the hypomorphic and LKB1 knockout muscles, contraction induced normal T-loop phosphorylation of the ERK1/ERK2 kinases (Figure 4B), which are potently activated by contraction or exercise (Sakamoto and Goodyear, 2002). Consistent with previous reports in rat muscle (Sakamoto *et al*, 2004), LKB1 activity in mouse muscle was not affected by contraction (Figure 4D).

LKB1 is required for contraction-induced glucose transport in skeletal muscle

Muscle contraction stimulates glucose uptake, but the contribution that AMPK makes to this process relative to other signalling pathways has been controversial (Aschenbach




Figure 4 Role of LKB1 in regulating contraction-induced AMPK activation. (A) One leg from anaesthetised mice of the indicated genotype (7–10 weeks of age) was subjected to *in situ* hindlimb muscle contraction (Ctr, contraction) via sciatic nerve stimulation for 5 min, and the other leg served as noncontracted control (B, basal). Tibialis anterior and EDL muscles from both legs were rapidly extracted and snap frozen in liquid nitrogen. AMPK α 2 was immunoprecipitated and assayed with the AMARA peptide. Assays were performed in duplicate from muscles derived from four to five mice and results are presented as the average \pm s.e.m. (B) Equal amounts of protein (20–30 μ g) from the muscle extracts were immunoblotted with the indicated antibodies. Immunoblot analysis of ACC phosphorylation was also assessed by quantitative Li-Cor analysis. The immunoblotting results are representative of independent experiments performed with tissues from at least three mice. (C, D) As in (A), except that AMPK α 1 (C) or LKB1 (D) was immunoprecipitated and assayed with AMARA or LKBtide. * P < 0.05 basal versus contraction within each genotype; (a) P < 0.05 versus LKB1^{+/+} (basal); (b) P < 0.05 versus LKB1 ^{β / β} Cre^{-/-} (basal) (c) P < 0.05 versus LKB1^{+/+} (ctr); (d) P < 0.05 versus LKB1 ^{β / β} Cre^{-/-} (ctr). Statistical analysis performed using one-way ANOVA and Tukey's *post hoc* test.

et al, 2004). We therefore measured glucose uptake in EDL muscle using *in situ* (Figure 5A) as well *in vitro* (Figure 5B) methodologies to induce muscle contraction. For the *in situ* approach, contraction was induced through sciatic nerve stimulation for 5 min and then EDL muscles were isolated and glucose transport measured following immersion for 10 min in ³H-2-deoxyglucose. This revealed that, in LKB1^{+/+} or hypomorphic LKB1 ^{β / β} Cre^{-/-} muscle, basal levels of glucose transport were similar and contraction stimulated the uptake of ³H-2-deoxyglucose two-fold (Figure 5A). In contrast, in the LKB1-deficient LKB1 ^{β / β} Cre^{+/-} muscle, the basal uptake of ³H-2-deoxyglucose, although similar to that observed in LKB1-expressing muscle, was marginally stimu-

lated, even though this was not statistically significant (Figure 5A). In the *in vitro* approach, isolated EDL muscles were induced to contract for 10 min by electrical stimulation and then glucose transport quantified following immersion of the muscle for 10 min in ³H-2-deoxyglucose. This revealed that the 1.7-fold stimulation of glucose uptake induced by contraction in wild-type and hypomorphic muscles was largely blocked in the LKB1-deficient LKB1 ^{β / β} Cre^{+/-} muscle (Figure 5B). We also monitored muscle force production during *in vitro* contraction and observed that, in the LKB1-lacking or LKB1-hypomorphic muscles, the force profiles were identical to those of wild-type muscle, indicating that the ability of LKB1-deficient muscles to contract is not




Figure 5 Role of LKB1 in regulating contraction-induced glucose transport. (A) One leg from anaesthetised mice of the indicated genotype (7–10 weeks of age) was subjected to *in situ* hindlimb muscle contraction (Ctr, contraction) via sciatic nerve stimulation for 5 min, and the other leg served as noncontracted control (B, basal). Following contraction, EDL muscle was isolated and glucose transport measured as described in Materials and methods. (B) Isolated EDL muscles were either subjected to *in vitro* contraction (Ctr, contraction) via electrical stimulation for 10 min or left unstimulated (B, basal). Glucose transport was then measured as described in Materials and methods. The data in (A) and (B) are presented as the mean \pm s.e.m. for muscle isolated from the indicated number (*n*) of mice of each genotype. (C) Force production during contraction protocol. Isolated EDL muscles were subjected to 10 sets of 10-s *in vitro* contraction and total force generated for each contraction was calculated as described in Supplementary data. The data are presented as the mean \pm s.e.m. for muscle isolated from three mice for each genotype. (D) Glucose transport activity in isolated EDL muscle from the indicated mice, determined without (–) or with 100 nM insulin (INS) as described in Materials and methods. The data are presented as the mean \pm s.e.m. for muscle isolated from three mice for each genotype. **P* < 0.05 basal versus contraction within each genotype; #*P* < 0.05 versus LKB1^{+/+} (contraction). Statistical analysis performed using one-way ANOVA and Tukey's *post hoc* test.

impaired (Figure 5C). As a further control to ensure that the glucose transport apparatus was functional in the LKB1-deficient muscle, we also measured glucose transport in the presence or absence of insulin, which stimulates the process through an AMPK-independent pathway. This revealed that insulin stimulated 2-deoxyglucose uptake two-fold in the LKB1-lacking LKB1^{fl/fl}Cre^{+/-} muscle, similar to that observed in LKB1-expressing muscle (Figure 5D).

LKB1 is required for phenformin-induced AMPK α 2 activation

To study whether LKB1 was required for the activation of AMPK α 2 by the antidiabetic drug phenformin, we treated isolated EDL muscle in the presence or absence of 2 mM phenformin for 60 min. Phenformin was employed rather than metformin, as this close relative of metformin activates AMPK much more potently in muscle than metformin (KS, unpublished results). Phenformin induced a three-fold stimulation of AMPK α 2 in LKB1-expressing muscle (Figure 6A), which was accompanied by a marked increase in phosphorylation of AMPK at Thr172 and ACC at Ser212 (Figure 6B). In the LKB1 hypomorphic muscle, the basal activity of AMPK α 2 was reduced, but phenformin still stimulated AMPK α 2, albeit to a level lower than observed in wild-type muscle

(Figure 6A). By contrast, in LKB1-deficient muscle, phenformin failed to significantly increase the low AMPK α 2 activity or the phosphorylation of AMPK at Thr172 and ACC at Ser212. We attempted to measure phenformin-stimulated glucose uptake in EDL muscles. However, under the conditions employed, we were unable to measure any significant increase in glucose uptake in LKB1-expressing muscles (KS, data not shown). To our knowledge, no other group has reported that phenformin can stimulate glucose uptake in isolated muscle.

Role of LKB1 in controlling cellular energy charge

Activation of AMPK during muscle contraction is thought to regulate cellular energy balance (Hardie *et al*, 2003). We therefore measured ADP:ATP and AMP:ATP ratios in control and contracted muscles (tibialis anterior and EDL) by quantitative capillary electrophoresis (Figure 7). As AMP can be converted to IMP by AMP deaminase, we also measured IMP:ATP ratios. In unstimulated LKB1-deficient LKB1^{fl/fl}Cre^{+/-} muscle, the ADP:ATP, AMP:ATP and IMP:ATP ratios were similar to those found in resting LKB1^{+/+} and LKB1^{fl/fl} muscle (Figure 7). In wild-type LKB1^{+/+} muscle, contraction only moderately increased the ADP:ATP (Figure 7A) and AMP:ATP (Figure 7B) ratios.




Figure 6 Role of LKB1 in regulating phenformin-induced AMPK α 2 activation. (A) Isolated mouse EDL muscles derived from wild-type and mutant mice (7–10 weeks of age) were incubated in the presence (Phen) or absence (–) of 2 mM phenformin for 60 min. AMPK α 2 was immunoprecipitated and assayed with the AMARA peptide. Assays were performed in duplicate from tissues derived from four mice and results are presented as the average \pm s.e.m. (B) Equal amounts of protein (20–30 μ g) from the muscle extracts were immunoblotted with the indicated antibodies. The immunoblotting results are representative of independent experiments performed with tissues from at least two mice. * P < 0.05 basal versus phenformin within each genotype; (a) P < 0.05 versus LKB1^{+/+} (basal); (b) P < 0.05 versus LKB1^{fl/fl}Cre^{-/-} (basal); (c) P < 0.05 versus LKB1^{+/+} (Phen); (d) P < 0.05 versus LKB1^{fl/fl}Cre^{-/-} (Phen). Statistical analysis performed using one-way ANOVA and Tukey's *post hoc* test.

However, in LKB1 hypomorphic muscle, contraction increased ADP:ATP and AMP:ATP ratios to statistically significant higher levels than those observed in wild-type muscle. These levels were further increased in the LKB1^{fl/fl}Cre^{+/-} muscle. In LKB1^{+/+} muscle, IMP:ATP ratios were similar to those observed in LKB1^{fl/fl}Cre^{+/-} muscle, in which the basal ratios were very low, but increased ~100-fold after contraction (Figure 7C). We found that in the LKB1 hypomorphic as well as the LKB1^{fl/fl}Cre^{+/-} muscles the IMP:ATP ratios were further increased by contraction.

Discussion

Our results establish that LKB1 is a major regulator of AMPK α 2 activity in skeletal muscle, and that lack of LKB1 is not compensated for by other kinases. Our results also support the notion that activation of AMPK plays a major role in maintaining cellular energy levels. This is based on the finding that, in LKB1-deficient muscles in which AMPK is not activated, muscle contraction resulted in an abnormal elevation of the ADP:ATP, AMP:ATP and IMP:ATP ratios (Figure 7). We believe that this is the first genetic evidence directly demonstrating the important role that LKB1 pathway plays




Figure 7 Measurement of ADP:ATP, AMP:ATP and IMP:ATP ratios in contracting muscle. One leg from anaesthetised mice of the indicated genotype (7–10 weeks of age) was subjected to *in situ* hindlimb muscle contraction (Ctr, contraction) via sciatic nerve stimulation for 5 min, and the other leg served as noncontracted control (B, basal). Following contraction, tibialis anterior and EDL muscle was isolated and nucleotides extracted and analysed by capillary electrophoresis as described in Supplementary data. The ratios of ADP:ATP (A), AMP:ATP (B) and IMP:ATP (C) derived from analysis of three independent muscle samples for each condition were measured. The results are shown as the average \pm s.e.m. * P < 0.05 basal versus contraction within each genotype; (a) P < 0.05 versus LKB1^{+/+} (basal); (b) P < 0.05 versus LKB1^{fl/fl}Cre^{-/-} (ctr). Statistical analysis performed using one-way ANOVA and Tukey's *post hoc* test.

in maintaining cellular energy balance *in vivo*. Moreover, our data clarify the mechanism by which muscle contraction activates AMPK. This is likely to be through elevation of AMP levels via the action of adenylate kinase, subsequent to depletion of ATP. AMP then interacts with the AMPK γ subunit, inducing conformational changes in the AMPK complex that stimulate the rate of phosphorylation of the AMPK α catalytic subunit by the LKB1 complex.

In contrast to the results with AMPK α 2, we observed that AICAR (Figure 3) as well as phenformin (data not shown) still induced substantial activation of AMPK α 1, in LKB1-deficient muscle. However, the finding that, in LKB1 hypomorphic muscle expressing 10-fold lower levels of LKB1, both the basal and AICAR-stimulated AMPK α 1 activity were reduced three- to four-fold (Figure 3) provides genetic evidence that LKB1 does regulate the activity of AMPK α 1. A likely reason as to why AMPK α 1 activity was not further reduced by expression of Cre recombinase could be that AMPK α 1 activity was derived mainly from nonmuscle cell types, such as fibroblasts, endothelial, fat or blood cells, which are known to be present in the muscle tissue and which would not express the Cre recombinase. This could also explain why AMPK α 1 is not activated in contracting muscle under conditions where AMPK α 2 was stimulated (Figure 4). However, our data do not rule out the possibility that AMPK α 1 is regulated by an LKB1-independent mechanism that is not triggered by contraction. LKB1 does activate AMPK complexes containing either AMPK α 1 or AMPK α 2 in cell-free assays (Hawley *et al*, 2003). Most studies in LKB1-deficient cell lines have been performed using approaches that immunoprecipitate both AMPK α 1 and AMPK α 2 activities (Hawley *et al*, 2003; Woods *et al*, 2003b; Shaw *et al*, 2004). However, in one study in which AMPK α 1 activity was assayed specifically, AICAR and phenformin were unable to stimulate AMPK α 1 activity in LKB1-deficient HeLa cells and mouse embryo fibroblasts (Lizcano *et al*, 2004), demonstrating that LKB1 can regulate AMPK α 1 activity in intact cultured cell lines.

Our finding that male LKB1^{fl/fl} mice are infertile suggests a role for LKB1 in regulating spermatogenesis and/or sperm motility. In testis, a faster migrating 48 kDa species of LKB1 is detected by antibodies raised against an N-terminal peptide of LKB1, which is absent in the LKB1^{fl/fl} mice (Figure 2F). In the LKB1^{fl/fl} mice, LKB1 is expressed after exon 4 from a cDNA fragment, therefore suggesting that the 48 kDa species is formed as a C-terminal splice variant of LKB1. In future work, it will be of interest to explore whether the faster migrating form of LKB1 plays a specific role in testis, and whether AMPK or one of the other AMPK-related kinases activated by LKB1 play a role in spermatogenesis and/or sperm motility. The SNRK kinase that is activated by LKB1 is specifically expressed in testis (Jaleel *et al*, 2005).

Our data demonstrate that a 10-fold reduction in LKB1 expression in the muscle of LKB1^{fl/fl} mice reduced basal AMPK α 2 activity as well as its activation by AICAR and contraction, although by much less than 10-fold. This suggests that, in muscle tissues, the normal level of LKB1 is not rate-limiting and exerts only a small degree of control over the activation of AMPK α 2. This may be a common feature of protein kinase signalling networks, as we have previously shown that, in hypomorphic mice that express only 10% of the normal level of PDK1, two of its downstream targets, that is, protein kinase B and ribosomal protein S6 kinase, are activated normally (Lawlor *et al*, 2002). Our results also illustrate the sensitivity and spare capacity that exists in the LKB1–AMPK signalling pathway. For example, AMPK α 2 activation induced by contraction was reduced ~2-fold in the hypomorphic LKB1^{fl/fl} mice, but this had no significant effect on the downstream actions of AMPK that we have investigated, namely phosphorylation of ACC (Figure 4B) or stimulation of

glucose uptake (Figure 5A and B). Thus, only very small increases in AMPK activity may be required to stimulate these downstream processes, and above that level the system becomes saturated. We are unable to rule out the possibility that the LKB1 hypomorphic mice may have adapted in order to compensate for the reduction in expression of LKB1. Humans with Peutz–Jeghers syndrome lacking one allele of the *LKB1* gene have not been reported to suffer from an inability to exercise or to possess any metabolic disorders. Our findings that AMPK is significantly activated in LKB1 hypomorphic mice may suggest that a single copy of the LKB1 allele in humans with Peutz–Jeghers syndrome is sufficient to normally regulate most processes downstream of AMPK.

In LKB1-deficient muscle, we found that phosphorylation of ACC, although greatly reduced, was not abolished. The residual phosphorylation of ACC that is observed could be mediated by trace residual activation of AMPK α 2 in the LKB1-deficient muscle, or by a kinase other than AMPK that is also stimulated by contraction. We also noticed that, in the hypomorphic mice, AICAR-induced, but not contraction-induced, phosphorylation of ACC was inhibited. This may provide further evidence that exercise can stimulate ACC phosphorylation independently of AMPK, and moreover it is also possible that exercise might promote ACC phosphorylation through the inhibition of a protein phosphatase. Interestingly, in LKB1-deficient fibroblasts, AICAR- and hydrogen-peroxide-induced phosphorylation of ACC was reduced, but not abolished (Shaw *et al*, 2004). Our findings suggest that AMPK is the major kinase phosphorylating the regulatory site (Ser212) on muscle ACC, but indicate that there may be other kinases that phosphorylate this site *in vivo*.

A complication that arises when generating muscle-specific knockouts using the Cre/loxP system is that it is sometimes difficult to obtain a complete ablation of gene expression throughout a muscle fibre, resulting from muscles being formed from fused myoblasts. Thus, if the Cre recombinase fails to delete the loxP-targeted gene in a small percentage of nuclei within a muscle fibre, this would result (in contrast to other tissues) in the targeted gene being expressed at low levels throughout the fibre. We have previously attempted to ablate PDK1 expression in skeletal muscle using the Cre/loxP system and found that, despite a 95% reduction in PDK1 expression in muscle tissues, the remaining PDK1 that is expressed throughout the muscle fibre is sufficient to normally activate its downstream substrates (Mora *et al*, 2003). In this study, we may have overcome the inherent problem of creating conditional knockouts in mouse muscle, by ablating LKB1 expression in a hypomorphic background. Had the LKB1^{fl/fl} mice not been hypomorphic for LKB1 expression, it is possible that a 95% reduction in LKB1 expression in the muscle of these animals would have only partially reduced AMPK α 2 activation and would not have affected contraction-induced ACC phosphorylation or glucose uptake.

A key finding of this study is that contraction-induced glucose uptake is largely inhibited in LKB1-deficient EDL muscle (Figure 5). This demonstrates that LKB1 is an important regulator of muscle contraction-induced glucose uptake. A previous study overexpressing dominant-negative AMPK from a muscle creatine kinase promoter (Mu *et al*, 2001) reported only a partial inhibition of glucose uptake, suggesting that AMPK-independent pathway(s) also regulate

contraction-induced glucose uptake. Recently, it was also reported that global knockouts of either AMPK α 1 or AMPK α 2 in mice had no effect on the ability of muscle contraction to induce glucose uptake (Jorgensen *et al*, 2004). By contrast, AICAR-induced glucose uptake was entirely blocked in the mice expressing the dominant-negative AMPK mutant in muscle (Mu *et al*, 2001) and in the AMPK α 2 knockout mice (Jorgensen *et al*, 2004). Our results with the LKB1 hypomorphic muscles, in which AMPK activation induced by contraction is considerably reduced, but glucose uptake is normal, indicate that a small activation of AMPK may be sufficient to induce normal glucose uptake. It is therefore possible that, in the muscles expressing the dominant-negative mutant, a small residual activation of endogenous AMPK is sufficient to cause a partial stimulation of glucose uptake. Similarly, in the individual AMPK α 2 knockout mice, there is a compensatory increase in the expression of AMPK α 1 (Viollet *et al*, 2003), and this might be sufficient to stimulate glucose uptake in the muscle. Interestingly, we also observed a two-fold increase in AMPK α 1 levels in the LKB1-deficient muscle, but presumably this would not be activated due to the lack of LKB1 and thus could not stimulate glucose uptake (Figure 3B). These data suggest that there is a negative feedback pathway by which expression of AMPK α 1 protein is upregulated following downregulation of AMPK activity, via an as yet undetermined mechanism. In contrast to the AMPK α 2 knockout mice (Viollet *et al*, 2003), the LKB1-deficient animals are not glucose intolerant. A likely explanation for this difference is that glucose intolerance in the AMPK α 2-deficient mice arises from defects in the AMPK kinase signalling pathways in tissues such as the autonomic nervous system, rather than in muscle (Viollet *et al*, 2003).

Our results do not exclude the possibility that LKB1 may be able to stimulate contraction-induced glucose uptake through an AMPK-independent mechanism. The most likely candidates for downstream targets of LKB1 that might mediate this effect would be the AMPK-related kinases that are also phosphorylated and activated by LKB1 (Lizcano *et al*, 2004). Although we were previously unable to demonstrate that any of these enzymes that were detectable in skeletal muscle (QSK, QIK, MARK2, MARK3 and MARK4) were stimulated by muscle contraction, AICAR or phenformin (Sakamoto *et al*, 2004), we cannot rule out the possibility that these enzymes might play a permissive role in enabling contraction to induce glucose uptake.

In conclusion, we provide here the first genetic evidence of the crucial role that LKB1 plays in regulating AMPK activation in muscle. Our data suggest that contraction/exercise-induced activation of AMPK and glucose uptake requires LKB1. We also demonstrate for the first time that contraction in muscles lacking AMPK activity results in abnormal increases in cellular AMP:ATP ratio, supporting the notion that LKB1 network functions as a sensor and regulator of cellular energy charge that protects muscle cells against the metabolic stress of contraction.

Materials and methods

Materials and details of antibodies and procedures used to prepare tissue lysates, perform immunoblotting, assay of LKB1/AMPK as well as glucose transport, contraction force, nucleotide measurements and statistics employed are described in Supplementary data.

Mice breeding and genotype analysis

All animal studies and breeding were approved by the University of Dundee ethical committee and performed under a UK Home Office project license. LKB1^{fl/fl} mice were crossed to transgenic mice expressing Cre recombinase under muscular creatine kinase promoter (Cre) (Bruning *et al*, 1998), which had been backcrossed for seven generations to the C57BL/6J strain. Genotyping was performed by PCR using genomic DNA isolated from tails. The presence of a wild-type or floxed LKB1 allele was detected using two primers, p200, 5'-CCAGCCTTCTGACTCTCAGG-3' and p201, 5'-GTAGGTATTCCAGGCCGTC-3'. For the detection of CRE, the following primers were employed: Cre1, 5'-AAATGGTTCCCGCA GAACC-3' and Cre10, 5'-TAGCTGGCTGGTGGCAGATG-3'.

Incubation of isolated muscle with AICAR or phenformin

Mice (7–10 weeks of age) were fasted overnight (16 h) prior to experiment and killed by cervical dislocation, and EDL muscles were rapidly and carefully removed. Tendons from both ends of each muscle were tied with suture (silk 4–0) and mounted on an incubation apparatus. The muscles were incubated as described previously (Sakamoto *et al*, 2004). Muscles were incubated in 8 ml of Krebs-Ringer bicarbonate (KRB) buffer (117 mM NaCl, 2.5 mM CaCl₂, 1.2 mM KH₂PO₄, 1.2 mM MgSO₄, 24.6 mM NaHCO₃, pH 7.4) containing 2 mM pyruvate for 60 min at 37°C, in the presence or absence of 2 mM AICAR or 2 mM phenformin. At the end of the incubation period, muscles were quickly frozen in liquid nitrogen. Muscles were stored at –80°C.

In situ muscle contraction

Mice (7–10 weeks of age) were fasted overnight (16 h) and were anaesthetised with sodium pentobarbital (90 mg/kg of body weight, administered intraperitoneally); the sciatic nerves to both legs were surgically exposed, and electrodes were attached. One leg was subjected to electrical stimulation for 5 min (train rate, 1/s; train duration, 500 ms; pulse rate, 100 Hz; duration, 0.1 ms at 2–5 V), and the other leg served as sham-operated (noncontracted) control as described previously (Sakamoto *et al*, 2002). Immediately after nerve stimulation, mice were killed by cervical dislocation, and tibialis anterior and EDL muscles were rapidly removed and then frozen in liquid nitrogen. Muscle tissues were stored at –80°C.

In vitro muscle contraction

Mice (7–10 weeks of age) were fasted overnight (16 h) prior to experiment and killed by cervical dislocation, EDL muscles were rapidly removed. Tendons from both ends of each muscle were tied with suture and mounted on an incubation apparatus with resting length. The muscles were incubated in 8 ml of KRB buffer containing 2 mM pyruvate for 50 min at 37°C. When contracted, muscles were electrically stimulated during the last 10 min of this period (train rate, 2/min; train duration, 10 s; pulse rate, 100 Hz; duration, 0.1 ms at 100 V).

Supplementary data

Supplementary data are available at *The EMBO Journal* Online.

Acknowledgements

We thank Ronald Kahn (Joslin Diabetes Center) for providing the Cre expressing mice, Gail Fraser, Victoria Murray-Tait and Laura Armit for technical assistance, Jose Bayascas and Kirsty Mustard for helpful advice, Greg Stewart and the protein production and antibody purification teams (Division of Signal Transduction Therapy (DSTT), University of Dundee) co-ordinated by Hilary McLauchlan and James Hastie for affinity purification of antibodies, the Sequencing Service (School of Life Sciences, University of Dundee, Scotland, www.dnaseq.co.uk) for DNA sequencing. We thank the Association for International Cancer Research (DRA), Cancer Research UK and Breakthrough Breast Cancer (AA), Diabetes UK (DRA+DG), the Medical Research Council UK (DRA+DG), the Moffat Charitable Trust, the European Commission (DGH, QLGI-CT-2001-0148800 RTD contract), the Wellcome Trust (DGH), as well as the pharmaceutical companies that support the Division of Signal Transduction Therapy (AstraZeneca, Boehringer-Ingelheim, GlaxoSmithKline, Merck & Co. Inc., Merck KGaA and Pfizer) for financial support.

References

- Altarejos JY, Taniguchi M, Clanachan AS, Lopaschuk GD (2005) Myocardial ischemia differentially regulates LKB1 and an alternate 5'AMP-activated protein kinase kinase. *J Biol Chem* **280**: 183–190
- Aschenbach WG, Sakamoto K, Goodyear LJ (2004) 5' adenosine monophosphate-activated protein kinase, metabolism and exercise. *Sports Med* **34**: 91–103
- Baas AF, Boudeau J, Sapkota GP, Smit L, Medema R, Morrice NA, Alessi DR, Clevers HC (2003) Activation of the tumour suppressor kinase LKB1 by the STE20-like pseudokinase STRAD. *EMBO J* **22**: 3062–3072
- Baron SJ, Li J, Russell III RR, Neumann D, Miller EJ, Tuerk R, Wallimann T, Hurley RL, Witters LA, Young LH (2005) Dual mechanisms regulating AMPK kinase action in the ischemic heart. *Circ Res* **96**: 337–345
- Boudeau J, Baas AF, Deak M, Morrice NA, Kieloch A, Schutkowski M, Prescott AR, Clevers HC, Alessi DR (2003) MO25alpha/beta interact with STRADalpha/beta enhancing their ability to bind, activate and localize LKB1 in the cytoplasm. *EMBO J* **22**: 5102–5114
- Boudeau J, Scott JW, Resta N, Deak M, Kieloch A, Komander D, Hardie DG, Prescott AR, van Aalten DM, Alessi DR (2004) Analysis of the LKB1–STRAD–MO25 complex. *J Cell Sci* **117**: 6365–6375
- Bruning JC, Michael MD, Winnay JN, Hayashi T, Horsch D, Accili D, Goodyear LJ, Kahn CR (1998) A muscle-specific insulin receptor knockout exhibits features of the metabolic syndrome of NIDDM without altering glucose tolerance. *Mol Cell* **2**: 559–569
- Carling D (2004) The AMP-activated protein kinase cascade—a unifying system for energy control. *Trends Biochem Sci* **29**: 18–24
- Corton JM, Gillespie JG, Hawley SA, Hardie DG (1995) 5-aminimidazole-4-carboxamide ribonucleoside. A specific method for activating AMP-activated protein kinase in intact cells? *Eur J Biochem* **229**: 558–565
- Davies SP, Sim AT, Hardie DG (1990) Location and function of three sites phosphorylated on rat acetyl-CoA carboxylase by the AMP-activated protein kinase. *Eur J Biochem* **187**: 183–190
- Fryer LG, Fougelle F, Barnes K, Baldwin SA, Woods A, Carling D (2002) Characterization of the role of the AMP-activated protein kinase in the stimulation of glucose transport in skeletal muscle cells. *Biochem J* **363**: 167–174
- Fujii N, Hayashi T, Hirshman MF, Smith JT, Habinowski SA, Kaijser L, Mu J, Ljungqvist O, Birnbaum MJ, Witters LA, Thorell A, Goodyear LJ (2000) Exercise induces isoform-specific increase in 5'AMP-activated protein kinase activity in human skeletal muscle. *Biochem Biophys Res Commun* **273**: 1150–1155
- Hardie DG (2004) The AMP-activated protein kinase pathway—new players upstream and downstream. *J Cell Sci* **117**: 5479–5487
- Hardie DG, Scott JW, Pan DA, Hudson ER (2003) Management of cellular energy by the AMP-activated protein kinase system. *FEBS Lett* **546**: 113–120
- Hawley SA, Boudeau J, Reid JL, Mustard KJ, Udd L, Makela TP, Alessi DR, Hardie DG (2003) Complexes between the LKB1 tumor suppressor, STRADalpha/beta and MO25alpha/beta are upstream kinases in the AMP-activated protein kinase cascade. *J Biol* **2**: 28
- Hayashi T, Hirshman MF, Kurth EJ, Winder WW, Goodyear LJ (1998) Evidence for 5' AMP-activated protein kinase mediation of the effect of muscle contraction on glucose transport. *Diabetes* **47**: 1369–1373
- Hemminki A, Markie D, Tomlinson I, Avizienyte E, Roth S, Loukola A, Bignell G, Warren W, Aminoff M, Högglund P, Jarvinen H, Kristo P, Pelin K, Ridanpaa M, Salovaara R, Toro T, Bodmer W, Olschwang S, Olsen AS, Stratton MR, de la Chapelle A, Aaltonen LA (1998) A serine/threonine kinase gene defective in Peutz–Jeghers syndrome. *Nature* **391**: 184–187
- Hong SP, Leiper FC, Woods A, Carling D, Carlson M (2003) Activation of yeast Snf1 and mammalian AMP-activated protein kinase by upstream kinases. *Proc Natl Acad Sci USA* **100**: 8839–8843
- Jaleel M, McBride A, Lizcano JM, Deak M, Toth R, Morrice NA, Alessi DR (2005) Identification of the sucrose non-fermenting related kinase SNRK, as a novel LKB1 substrate. *FEBS Lett* **579**: 1417–1423
- Jenne DE, Reimann H, Nezu J, Friedel W, Löff S, Jeschke R, Müller O, Back W, Zimmer M (1998) Peutz–Jeghers syndrome is caused by mutations in a novel serine threonine kinase. *Nat Genet* **18**: 38–43
- Jorgensen SB, Viollet B, Andreelli F, Frosig C, Birk JB, Schjerling P, Vaulont S, Richter EA, Wojtaszewski JF (2004) Knockout of the alpha2 but not alpha1 5'-AMP-activated protein kinase isoform abolishes 5-aminimidazole-4-carboxamide-1-beta-4-ribofuranosidebut not contraction-induced glucose uptake in skeletal muscle. *J Biol Chem* **279**: 1070–1079
- Lawlor MA, Mora A, Ashby PR, Williams MR, Murray-Tait V, Malone L, Prescott AR, Lucocq JM, Alessi DR (2002) Essential role of PDK1 in regulating cell size and development in mice. *EMBO J* **21**: 3728–3738
- Lizcano JM, Goransson O, Toth R, Deak M, Morrice NA, Boudeau J, Hawley SA, Udd L, Makela TP, Hardie DG, Alessi DR (2004) LKB1 is a master kinase that activates 13 kinases of the AMPK subfamily, including MARK/PAR-1. *EMBO J* **23**: 833–843
- Merrill GF, Kurth EJ, Hardie DG, Winder WW (1997) AICA riboside increases AMP-activated protein kinase, fatty acid oxidation, and glucose uptake in rat muscle. *Am J Physiol* **273**: E1107–E1112
- Mora A, Davies AM, Bertrand L, Sharif I, Budas GR, Jovanovic S, Mouton V, Kahn CR, Lucocq JM, Gray GA, Jovanovic A, Alessi DR (2003) Deficiency of PDK1 in cardiac muscle results in heart failure and increased sensitivity to hypoxia. *EMBO J* **22**: 4666–4676
- Mu J, Brozinick Jr JT, Valladares O, Bucan M, Birnbaum MJ (2001) A role for AMP-activated protein kinase in contraction- and hypoxia-regulated glucose transport in skeletal muscle. *Mol Cells* **7**: 1085–1094
- Nath N, McCartney RR, Schmidt MC (2003) Yeast Pak1 kinase associates with and activates Snf1. *Mol Cell Biol* **23**: 3909–3917
- Sakamoto K, Goodyear LJ (2002) Invited review: intracellular signaling in contracting skeletal muscle. *J Appl Physiol* **93**: 369–383
- Sakamoto K, Goransson O, Hardie DG, Alessi DR (2004) Activity of LKB1 and AMPK-related kinases in skeletal muscle; effects of contraction, phenformin and AICAR. *Am J Physiol Endocrinol Metab* **287**: E310–E317
- Sakamoto K, Hirshman MF, Aschenbach WG, Goodyear LJ (2002) Contraction regulation of Akt in rat skeletal muscle. *J Biol Chem* **277**: 11910–11917
- Scott JW, Hawley SA, Green KA, Anis M, Stewart G, Scullion GA, Norman DG, Hardie DG (2004) CBS domains form energy-sensing modules whose binding of adenosine ligands is disrupted by disease mutations. *J Clin Invest* **113**: 274–284
- Shaw RJ, Kosmatka M, Bardeesy N, Hurley RL, Witters LA, DePinho RA, Cantley LC (2004) The tumor suppressor LKB1 kinase directly activates AMP-activated kinase and regulates apoptosis in response to energy stress. *Proc Natl Acad Sci USA* **101**: 3329–3335
- Sutherland CM, Hawley SA, McCartney RR, Leech A, Stark MJ, Schmidt MC, Hardie DG (2003) Elm1p is one of three upstream kinases for the *Saccharomyces cerevisiae* SNF1 complex. *Curr Biol* **13**: 1299–1305
- Suzuki A, Kusakai G, Kishimoto A, Shimojo Y, Ogura T, Lavin MF, Esumi H (2004) IGF-1 phosphorylates AMPK-alpha subunit in ATM-dependent and LKB1-independent manner. *Biochem Biophys Res Commun* **324**: 986–992
- Viollet B, Andreelli F, Jorgensen SB, Perrin C, Geloën A, Flamez D, Mu J, Lenzner C, Baud O, Bennoun M, Gomas E, Nicolas G, Wojtaszewski JF, Kahn A, Carling D, Schuit FC, Birnbaum MJ, Richter EA, Burcelin R, Vaulont S (2003) The AMP-activated protein kinase alpha2 catalytic subunit controls whole-body insulin sensitivity. *J Clin Invest* **111**: 91–98
- Winder WW, Hardie DG (1999) AMP-activated protein kinase, a metabolic master switch: possible roles in type 2 diabetes. *Am J Physiol* **277**: E1–E10
- Woods A, Johnstone SR, Dickerson K, Leiper FC, Fryer LG, Neumann D, Schlattner U, Wallimann T, Carlson M, Carling D (2003a) LKB1 is the upstream kinase in the AMP-activated protein kinase cascade. *Curr Biol* **13**: 2004–2008
- Woods A, Vertommen D, Neumann D, Turk R, Bayliss J, Schlattner U, Wallimann T, Carling D, Rider MH (2003b) Identification of phosphorylation sites in AMP-activated protein kinase (AMPK) for upstream AMPK kinases and study of their roles by site-directed mutagenesis. *J Biol Chem* **278**: 28434–28442
- Zhou G, Myers R, Li Y, Chen Y, Shen X, Fenyk-Melody J, Wu M, Ventre J, Doebber T, Fujii N, Musi N, Hirshman MF, Goodyear LJ, Moller DE (2001) Role of AMP-activated protein kinase in mechanism of metformin action. *J Clin Invest* **108**: 1167–1174